

HPB Moorhampton Legume trial 1992-3

White clover establishment with nurse species

Tree girths at end of first season[Ashton Bitter]

| | row 1 | | row 2 | | row 3 | | |
|----------------------|-------|-------|-------|-------|-------|-------|---------------|
| TREATMENT | rep 1 | rep 2 | rep 1 | rep 2 | rep 1 | rep 2 | Means |
| Bare | 22.0 | 22.0 | 23.5 | 22.5 | 21.5 | 21.5 | 22.2 |
| S184 | 20.0 | 21.0 | 22.0 | 24.5 | 21.5 | 22.0 | 21.8 |
| S184+fescue/mustard | 25.0 | 23.0 | 24.5 | 24.0 | 23.0 | 22.0 | 23.6 * |
| S184+legume/ryegrass | 21.5 | 21.5 | 21.5 | 22.5 | 22.0 | 22.5 | 21.9 |
| Kent | 21.0 | 23.5 | 22.0 | 20.5 | 22.0 | 22.0 | 21.8 |
| Kent+fescue/mustard | 25.5 | 22.0 | 22.0 | 23.5 | 22.5 | 23.0 | 23.1 * |
| Kent+legume/ryegrass | 22.5 | 24.0 | 22.5 | 22.0 | 21.5 | 21.5 | 22.3 |
| Huia | 24.0 | 23.0 | 22.5 | 17.0 | 20.5 | 21.5 | 21.4 |
| Huia+fescue/mustard | 23.0 | 21.0 | 22.5 | 22.0 | 22.5 | 22.5 | 22.3 |
| Huia+legume/ryegrass | 22.0 | 18.0 | 22.5 | 23.0 | 21.0 | 22.0 | 21.4 |
| | | | | | | | |
| ROW/REP MEANS | 22.7 | 21.9 | 22.6 | 22.2 | 21.8 | 22.1 | 22.2 |